


North Glasgow Hospitals Department of Haematology Service Users Handbook



Contents

Please note this document uses hyperlinks please “ctrl + click” on the “section” links in the table of contents and within text to move to the relevant section.

1.	Introduction	5
2.	General Information	5
2.1.	Regulation and Accreditation	5
2.2.	Complaints	6
2.3.	Result Enquiries	6
3.	Laboratory Hours	6
3.1.	Gartnavel General Hospital	6
3.2.	Glasgow Royal Infirmary	6
3.3.	Stobhill ACH	6
3.4.	West Glasgow ACH	6
3.5.	24 Hour Service	6
4.	Contact Details	7
4.1.	Postal Addresses	7
4.2.	Website	7
4.3.	Telephone Numbers	7
4.3.1.	Result Enquiries	8
4.3.2.	Gartnavel General Hospital Laboratory	8
4.3.3.	Glasgow Royal Infirmary Laboratory	8
4.3.4.	Stobhill ACH Laboratory	8
4.3.5.	West Glasgow ACH Laboratory	8
4.3.6.	Gartnavel General Hospital Clinical Staff	9
4.3.7.	Glasgow Royal Infirmary Clinical Staff	9
4.3.8.	Registrars (Glasgow Royal Infirmary)	10
4.3.9.	Registrars (Gartnavel General Hospital)	10
4.3.10.	Stobhill ACH Clinical Staff	10
4.3.11.	Senior Laboratory Staff (All sites)	10
4.3.12.	Lead Scientific Staff Gartnavel General Hospital	11
4.3.13.	Lead Scientific Staff Glasgow Royal Infirmary	11
4.3.14.	Transfusion Practitioner	12
5.	Urgent Samples, Advice and Result Interpretation	12
5.1.	Urgent Samples	12
5.1.1.	Glasgow Royal Infirmary and Gartnavel General Hospital	12
5.1.2.	Stobhill ACH	13
5.1.3.	West Glasgow ACH	13
5.1.4.	North East Glasgow GP's	13
5.1.5.	North West Glasgow GP's	13
5.2.	Advice	13
5.2.1.	North West Glasgow GP's	13
6.	Specimen Collection	14
6.1.	Specimen Type	14
6.2.	Sample Labelling	15
6.2.1.	Blood Transfusion Sample Labelling Requirements	15
6.2.2.	Consent	16
6.3.	Transportation of Samples	16
6.3.1.	Portering Services	16
6.3.2.	Vacuum Tube Specimen Delivery System	16
6.3.3.	Primary Care Specimen Collection Service	16
6.3.4.	Sending Specimens by Post	16

6.4. Restricted Specimens.....	17
7. Assay Repertoire and Turn Around Times.....	17
7.1. Haematology.....	17
7.1.1. Malaria Parasite Screening Request Requirements.....	17
7.1.2. Haemoglobinopathy Screening Request Requirements.....	18
7.2. Haemostasis.....	18
7.2.1. Haemostasis Assays General Requirements.....	19
7.2.2. Anticoagulation Therapy.....	20
7.2.3. Anti Xa Assays Special Requirements.....	20
7.2.4. Lupus Anticoagulant Assays Special Requirements.....	20
7.2.5. Anticoagulation Service.....	20
7.3. Haemato-oncology.....	20
7.3.1. Haemato-oncology sampling requirements.....	20
7.4. Blood Transfusion.....	21
7.4.1. Routine Blood Product Orders.....	21
7.4.2. Emergency Cross Match Requests.....	21
7.4.3. Atypical Red Cell Antibodies.....	22
7.4.4. Transfusion Reactions.....	22
7.4.5. Blood Components.....	22
7.5. Costs.....	23
8. Referred Assays.....	23
9. Reference Ranges.....	24
9.1. Haematology Reference Ranges.....	24
9.1.1. Haematology Reference Ranges for Adults.....	25
9.1.1. Haematology Reference Ranges for Children.....	25
9.1.2. Haematology Reference Ranges for Infants.....	26
9.1.3. ESR.....	26
9.2. Haemoglobinopathy Assays.....	27
9.3. Haemato-Oncology.....	27
9.3.1. Immunophenotyping.....	27
9.4. Haemostasis Reference Ranges.....	27
9.4.1. Routine Haemostasis Ranges.....	27
9.4.2. Specific Haemostasis Assay Reference Ranges.....	28
9.5. Haematinics Assays.....	29
9.6. Uncertainty of Measurement (UoM).....	29
9.6.1. Haematology UoM.....	30
9.6.2. Manual Leucocyte Differential UoM.....	31
9.6.3. Malaria Parasite Parasitaemia UoM.....	31
9.6.4. Bone Marrow Myelogram.....	31
9.6.5. Haemoglobinopathy UoM.....	31
9.6.6. Routine Haemostasis UoM.....	32
9.6.7. Special Haemostasis UoM.....	32
9.6.8. Haematinics Assays UoM.....	34
9.6.9. Haemato-Oncology UoM.....	34
9.6.10. Qualitative Assays UoM.....	34
9.7. Factors That Will Affect the Accuracy of Results.....	34
9.7.1. Common Factors.....	35
9.7.2. Blood Transfusion.....	35
9.7.3. Haemostasis.....	35
9.7.4. Haematology.....	36
9.7.5. Haemato-Oncology.....	36

Author	K Marriott	Authorised By	C McKie
--------	------------	---------------	---------

10. Confidentiality.....	36
11. Retention of Records	36
12. References	36
Appendix 1	38
Changes from Previous Version	38



1. Introduction

The NHSGG&C Diagnostics Division, Department of Haematology, North Sector provides a comprehensive routine and specialist haematology service. This service is provided by laboratories located at Gartnavel General Hospital, Glasgow Royal Infirmary, Stobhill Ambulatory Care Hospital and the West Glasgow Ambulatory Care Hospital.

The Department complies with national and international standards assessed by UKAS ISO 15189 and regulated by the MHRA and is committed to meeting the needs and requirements of service users.

This Handbook is designed to provide information about using the Haematology and Blood Transfusion service. Within this handbook, along with contact details can be found information about specimen requirements, specimen identification, request form requirements, safety considerations, transport, reference ranges, turnaround times and other information about assays as is appropriate.

2. General Information

The department currently provides services at four sites. These are Gartnavel General Hospital, Glasgow Royal Infirmary, Stobhill ACH and West Glasgow ACH

The Department provides a diagnostic laboratory service which includes:

- Core diagnostic laboratory services.
- Consultant led clinical advice and test interpretation.
- Regional specialised thrombosis and haemostasis laboratory service (Glasgow Royal Infirmary).
- A National Allogeneic Stem Cell and Regional Autologous Stem Cell processing service in association with the West of Scotland Bone Marrow Transplant Unit (Gartnavel General Hospital).
- Regional Immunophenotyping (cell marker) service for the diagnosis of haematological abnormalities and malignancies (Gartnavel General Hospital).
- Blood Transfusion Services (Glasgow Royal Infirmary and Gartnavel General Hospital).

Our Quality manual, quality policy and other information can be found on our webpage on the NHSGGC website please follow the link in section 4.2

2.1. Regulation and Accreditation

The Haematology Department for the North Glasgow Sector of Greater Glasgow and Clyde is regulated by the Medicines and Healthcare Products Regulatory Agency (MHRA) for compliance to The Medicines for Human Use (Clinical Trials) Regulations 2004 and the Blood Safety Quality Regulations 2005 (Amendment 2007), compliance with the Human Tissues Act 2004 by the Human Tissue Authority (HTA), compliance to the JACIE standards by The Joint Accreditation Committee ISCT-Europe and EBMT. It is also accredited by the United Kingdom Accreditation service (UKAS Number 9570) to the international standard ISO15189. Further information on the department's scope of accreditation to ISO15189 and accredited processes are available on the UKAS website or via the departments website. The department's schedule of scope can be found by following the link below.

[9570-Medical-Multiple.pdf \(ukas.com\)](https://www.ukas.com/9570-Medical-Multiple.pdf)

2.2. Complaints

Should there be cause to raise a complaint or users wish to feedback about the laboratory service a copy of the department’s feedback and complaints procedure can be found on the laboratory web page or from the Quality Manager on request. A copy of the NHSGGC complaints policy can be found on the NHSGGC website. A complaints form can be found on the department’s web page.

2.3. Result Enquiries

Telephoning for results can be wasteful of time, both in the clinical areas and in the Haematology department, before telephoning the department service users including primary care should check for results on Clinical Portal, TrakCare or any other relevant information systems. This is the fastest and most efficient way of obtaining results. If results are not available it is most likely that the analyses are not complete. Please only contact the laboratory directly for urgent results. All extremely abnormal results will be phoned to the requesting clinician, GP or clinical area as soon as they become available.

3. Laboratory Hours

3.1. Gartnavel General Hospital

Haematology and Haemostasis:	08:00 to 18:00 Monday to Friday
Blood Transfusion:	08:00 to 18:00 Monday to Friday
Haemato-Oncology:	09:00 to 17:00 Monday to Friday

The Blood Transfusion, Haematology and Haemostasis laboratory service is provided from Glasgow Royal Infirmary outside these hours.

3.2. Glasgow Royal Infirmary

24 Hour service provided (core hours 08:00 to 17:00)

3.3. Stobhill ACH

Haematology and Coagulation:	09:00 to 17:00 Monday to Friday
------------------------------	---------------------------------

The laboratory service is provided from Glasgow Royal Infirmary outside these hours

3.4. West Glasgow ACH

Haematology:	09:00 to 13:00 Wednesday ONLY .
--------------	--

This laboratory service is for the provision of the Haematology Clinic **ONLY**.

3.5. 24 Hour Service

The 24 hour laboratory service is provided from the Glasgow Royal Infirmary Site and consists primarily of a core haematology, core Haemostasis and blood transfusion service. Specific tests can be arranged by discussion with the on-call haematologist. There is no 24 hour service on site at Gartnavel General Hospital, Stobhill ACH or the West Glasgow ACH, this service is provided from the Glasgow Royal Infirmary Site.

4. Contact Details

4.1. Postal Addresses

Gartnavel General Hospital

Department of Haematology or Haemato-Oncology
 Gartnavel General Hospital,
 Paul O’Gorman Building
 21 Shelley Road
 Glasgow
 G12 0XB

Glasgow Royal Infirmary

Department of Haematology
 McEwan Building
 Glasgow Royal Infirmary
 Castle Street
 G4 0SF

Stobhill ACH

Haematology Laboratory
 Stobhill Ambulatory Care Hospital
 Stobhill
 Glasgow
 G21 3EW

West Glasgow ACH

Haematology Laboratory
 West Glasgow Ambulatory Care Hospital
 Yorkhill
 Glasgow
 G3 8SJ

4.2. Website

<https://www.nhsggc.scot/staff-recruitment/staff-resources/laboratory-medicine/haematology-and-blood-transfusion/>

4.3. Telephone Numbers

Please note some clinical and laboratory staff work at, or cover more than one site within the North Glasgow Sector.

4.3.1. Result Enquiries

Telephoning for results can be wasteful of time, both in the clinical areas and in the Haematology department, before telephoning the department service users including primary care should check for results on Clinical Portal, TrakCare or any other relevant information systems. This is the fastest and most efficient way of obtaining results. If results are not available it is most likely that the analyses are not complete. Please only contact the laboratory directly for urgent results.

All extremely abnormal results will be phoned to the requesting clinician, GP or clinical area as soon as they become available.

4.3.2. Gartnavel General Hospital Laboratory

External callers dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

Haematology and Haemostasis:	0141 301 Ext: (5)7721
Haemato-Oncology, Immunophenotyping and Stem Cell:	0141 301 Ext: (5)7708
Blood Transfusion:	0141 301 Ext: (5)7729

4.3.3. Glasgow Royal Infirmary Laboratory

External callers dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

Blood Transfusion	0141 242 Ext: (2)9603
	0141 242 Ext: (2)9604
	0141 242 Ext: (2)9606
Haemostasis	0141 242 Ext: (2)9605
Haematology	0141 242 Ext: (2)9601
Haematology	0141 242 Ext: (2)9602
Special Haemostasis	0141 242 Ext: (2)9552

4.3.4. Stobhill ACH Laboratory

External callers dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

All Laboratory Enquiries:	0141 355 Ext: (1)1469
---------------------------	-----------------------

4.3.5. West Glasgow ACH Laboratory

External callers dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

All Laboratory Enquiries:	0141 211 Ext: (8)6946
---------------------------	-----------------------

4.3.5.1. Out of Hours Contact Numbers

All Sites

Out of Hours (see laboratory hours in section 3 for details) the Glasgow Royal Infirmary Laboratory can be contacted by using the appropriate extension number. Clinical advice can be obtained by contacting switchboard and asking for the duty Haematologist.

There is no 24 hour service on site at Gartnavel General Hospital, Stobhill ACH or West Glasgow ACH, this services is provided from the Glasgow Royal Infirmary site.

4.3.6. Gartnavel General Hospital Clinical Staff

External callers dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

Consultants

Dr M Drummond 0141 301 Ext: (5)7734 email: mark.drummond@nhs.scot
 Secretary 0141 301 Ext: (5)7712

Dr M Leach 0141 301 Ext: (5)7736 email: mike.leach@nhs.scot
 Secretary 0141 301 Ext: (5)7713
 0141 301 Ext: (5)5305

Dr P McKay 0141 301 Ext: (5)7735 email: pam.mckay@nhs.scot
 Secretary 0141 301 Ext: (5)7711

Dr J Travers 0141 301 Ext: (5)7732 email: jennifer.travers@nhs.scot
 Secretary 0141 301 Ext: (5)7717

Dr C McDermott 0141 301 Ext: (5)7747 email: christopher.mcdermott@nhs.scot
 Secretary 0141 301 Ext: (5)7717

Dr R Soutar 0141 301 Ext: (5)7733 email: richard.soutar2@nhs.scot
 Secretary 0141 301 Ext: (5)7715

4.3.7. Glasgow Royal Infirmary Clinical Staff

Outside NHSGG&C, dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

Consultants:

Dr C Bagot 0141 242 Ext: (2)9594 email: catherine.bagot@nhs.scot
 Secretary 0141 201 Ext: (6)5306

Dr L McIlwaine 0141 242 Ext: (2)9595 email: louisa.mcilwaine@nhs.scot
 Secretary 0141 201 Ext: (6)5306

Dr R Rodgers 0141 242 Ext: (2)9593 email: ryan.rodgers@nhs.scot
 Secretary: 0141 201 Ext: (1)5305

Dr A Gibson Secretary	0141 242 Ext: (2)9554 0141 201 Ext: (1)3654	email: alison.gibson6@nhs.scot
Dr M Wilson Secretary	0141 242 Ext: (2)9592 0141 242 Ext: (1)3654	email: matthew.wilson@nhs.scot
Dr C Betts Secretary	0141 242 Ext (2)9504 0141 201 Ext (1)5305	email: caroline.betts2@nhs.scot
Dr S McNeill Secretary	0141 242 Ext (2)9583 0141 201 Ext (1)5305	email: susan.mcneill3@nhs.scot

4.3.8. Registrars (Glasgow Royal Infirmary)

On Duty Registrar	Page: 13733
Registrars' Office	0141 201 Ext: (1)3641 0141 201 Ext: (1)3655 0141 242 Ext: (2)9590 0141 242 Ext: (2)9591

4.3.9. Registrars (Gartnavel General Hospital)

Registrars' Office	0141 301 Ext: (5)7753
--------------------	-----------------------

4.3.10. Stobhill ACH Clinical Staff

External callers dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

Consultant:

Dr M Leach Secretary	0141 301 Ext: (5)7736 0141 301 Ext: (5)7713	email: mike.leach@nhs.scot
Dr L McIlwaine Secretary	0141 201 Ext: (1)3655 0141 201 Ext: (6)5306	email: louisa.mcilwaine@nhs.scot
Dr A Gibson Secretary	0141 242 Ext: (2)9554 0141 201 Ext: (1)3654	email: alison.gibson6@nhs.scot

4.3.11. Senior Laboratory Staff (All sites)

Outside NHSGG&C, dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

Technical Services Manager

Mrs Claire McKie	0141 242 Ext: (2)9529 (Glasgow Royal Infirmary) 0141 301 Ext: (5)7727 (Gartnavel General Hospital) 0141 355 Ext: (1)1469 (Stobhill ACH)
------------------	---

email: claire.mckie@nhs.scot

Sector Laboratory Manager

Ms Arlene David 0141 242 Ext: (2)9530 (Glasgow Royal Infirmary)
 0141 301 Ext: (5)7727 (Gartnavel General hospital)
 0141 355 Ext: (1)1469 (Stobhill ACH)
 email: arlene.david2@nhs.scot

Quality, Training and POCT Manager

Mr Kevin Marriott 0141 242 Ext: (2)9597 (Glasgow Royal Infirmary)
 0141 301 Ext: (5)7727 (Gartnavel General Hospital)
 0141 355 Ext: (1)1469 (Stobhill ACH)
 email: kevin.marriott@nhs.scot

4.3.12. Lead Scientific Staff Gartnavel General Hospital

Please note some laboratory staff work at, or cover more than one site within the North Glasgow Sector.

Outside NHS GG&C, dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

4.3.12.1. Blood Transfusion

Senior Biomedical Scientist 0141 301 Ext: (5)7727

4.3.12.2. Haemostasis

Senior Biomedical Scientist 0141 301 Ext: (5)7727

4.3.12.3. Haematology

Senior Biomedical Scientist 0141 301 Ext: (5)7727

4.3.12.4. Haemato-Oncology

Mrs Sharon Kelly 0141 301 Ext: (5)7707
 email: sharon.kelly4@nhs.scot

Consultant Clinical Scientist

Ms Gillian McGaffin 0141 301 Ext: (5)7709
 email: gillian.mcgaffin2@nhs.scot

4.3.13. Lead Scientific Staff Glasgow Royal Infirmary

Outside NHS GG&C, dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

4.3.13.1. Blood Transfusion

Ms Arlene David 0141 242 Ext: (2)9530
 email: arlene.david2@nhs.scot

Senior Biomedical Scientist 0141 242 Ext: (2)9541

4.3.13.2. Haemostasis

Mrs Caroline Lawrence 0141 242 Ext: (2)9596
 email: caroline.lawrence@nhs.scot

4.3.13.3. Haematology

Senior Biomedical Scientist 0141 242 Ext: (2)9539
 0141 242 Ext: (2)9540

4.3.14. Transfusion Practitioner

Outside NMSGG&C, dial full number omitting the number in brackets. Internal callers dial extension number prefixed by the number in brackets.

Glasgow Royal Infirmary and Stobhill ACH

Mrs Gillian O'Donnell 0770 2159063
 email: Gillian.odonnell2@nhs.scot

Gartnavel General Hospital

Mrs Tina Watson 0778 9616525
 Email: tina.watson2@nhs.scot

5. Urgent Samples, Advice and Result Interpretation

During normal hours, clinical and technical advice is available from scientific and clinical staff. Out with core hours clinical advice is available by contacting the duty haematologist via switchboard and technical advice is available by telephone contact of laboratory staff. Please note there is no Haematology service at Stobhill ACH after 17:00. There is no Haematology service after 18:00 or Haemato-Oncology service after 17:00 at Gartnavel General Hospital, all enquiries must then be directed to the GRI laboratory or on call clinical staff.

5.1. Urgent Samples

5.1.1. Glasgow Royal Infirmary and Gartnavel General Hospital

Please contact the relevant laboratory on either site using the telephone numbers in section 0. Please note there is no Haematology service after 18:00 and no Haemato-Oncology service after 17:00 at Gartnavel General Hospital.

5.1.2. Stobhill ACH

Please contact the Haematology laboratory on the telephone number in section 0 Please note as this is a rapid results service any sample received as routine or requiring a test not performed on the Stobhill ACH site may have already been transported to the GRI site, if advised that this has occurred please contact the GRI laboratory using the numbers in section 4.3.4 Please note there is no service at Stobhill ACH after 17:00

5.1.3. West Glasgow ACH

This provides a service to the Haematology clinic only. All other samples are processed at the Queen Elizabeth University Hospital and all enquiries must be directed there.

5.1.4. North East Glasgow GP's

If a result is required by 18:00 then please contact the Glasgow Royal Infirmary Laboratory using the number in section 4.3.6

5.1.5. North West Glasgow GP's

Samples are processed at the Queen Elizabeth University Hospital please contact the laboratory there to advise of urgent samples.

Please use the following telephone number for the QEUH laboratory: 0141 354 9105

5.2. Advice

Service Users can obtain technical advice from the laboratory including:

- Suitability of the assay
- Assays available in the laboratory.
- Significance of results
- Reference ranges

Please use the telephone numbers given in section 4.3 to contact the relevant scientific team or laboratory.

Service Users can obtain clinical advice including:

- Clinical suitability of the assay
- Treatment modality (if appropriate).
- Clinical interpretation of results

Please use the telephone numbers given in section 4.3.6 to contact the relevant clinical team.

5.2.1. North West Glasgow GP's

Samples for NW Glasgow primary care are processed at the Queen Elizabeth University Hospital (QEUH) however clinical advice is provided by the consultant team at Gartnavel general Hospital. To obtain clinical advice please use the contact numbers for the Gartnavel clinical team provided in section 4.3.6 for all other enquiries please contact the relevant QEUH laboratory.

6. Specimen Collection

The following information in this section and its subsections, applies at all times whether taking a single sample or multiple samples from an individual.

It is the responsibility of the staff undertaking sample collection to ensure that the correct sample bottle type is selected, labelled accurately, form completed with all relevant information and transported to the laboratory in a timely and correct manner.

The sample bottles must be checked to ensure that the bottles have not exceeded their expiry date. Samples received in bottles that have expired will be rejected as the results will not be accurate.

Please note the following points relevant to collection (or venepuncture) of good quality specimens (see also sections 7.1.1, 7.1.2, 7.2.1, 7.2.3, 7.2.4, 7.3.1 5.2):

- **CONFIRM THE IDENTITY** of the patient **PRIOR** to sampling
- **NEVER** pre-label specimen tubes
- Ideally, the patient must be resting for a full five minutes before specimen collection
- Use good quality veins
- **NEVER** take blood from a drip arm
- Do not take samples for Haemostasis studies from heparinised lines
- Avoid prolonged application of the tourniquet both for patient comfort and to avoid haemolysis within the specimen.
- Samples **MUST** be filled to the fill line as marked on the bottle. This is essential for Haemostasis assays.
- Following collection, specimen bottles containing anticoagulant must be inverted several times to ensure adequate mixing.
- **Do not** decant blood from one tube to another – there are different additives and these will give erroneous results.
- Following collection, ensure specimen bottle is labelled, as detailed in Section 6.2
- Ensure Request Form is completed, as detailed in Section 6.2 and that these details match those on the specimen bottle.
- Use a safe procedure at all times and dispose of sharps in sharps-boxes provided
- All specimens and request forms must be secured for transportation in the specimen compartment of an approved specimen transport bag
- Specimen tubes or request forms which are contaminated with blood will not be analysed
- Appropriate specimen containers must be used for each laboratory test.

6.1. Specimen Type

The appropriate containers for haematological tests are available from the stores on each site they are not supplied by the department. The type and amount of anticoagulant varies depending on the investigations requested and the amount of blood required.

A colour-coded specimen container (relating to type of anticoagulant and its use for individual laboratory tests) and vacuum assisted venepuncture system (Greiner Vacuette™) is used throughout NHS GG&C.

Wall charts and posters detailing the use of this system and the correct container for each test, are posted in most clinical areas throughout NHS GG&C, the specimen type is also detailed in sections 7.1 to 7.4. Additional guidance or information regarding the use of the Greiner Vacuette™ system may be obtained by contacting the Lead Phlebotomist.

A team of trained phlebotomists, covering all acute medical and surgical wards are available in all the North Sector Glasgow Hospitals. Public Holidays are covered by a limited service with details available from each Hospital Lead Phlebotomist.

6.2. Sample Labelling

Sample bottles **MUST** be labelled at the patient's side to avoid identification errors. **NEVER** place unlabelled samples in the same vicinity as others or label in a different area to the patient.

A fully completed request form must accompany a properly identified sample in all cases.

Minimal identifying particulars for Haematology and Haemostasis on **both** the request form and sample(s) (Not Blood Transfusion samples) are:

- Surname
- Forename
- CHI number
- Date of Birth

The form must also include:

- CHI number
- Gender
- Source of request i.e. ward and consultant in charge
- Brief clinical details
- Date of request
- Investigation requested
- Signature/name of requesting doctor and bleep number

Use of the Trackcare system will ensure that all of this required data will be present. Please note that due to interfacing limitations the assay requestor is not displayed in clinical portal, SCI store or IS.

Please note unlabelled or inadequately labelled samples **WILL NOT BE** accepted for analysis. In these circumstances the clinician or clinical area making the request will be notified and a fresh, suitably identified sample requested. Under **NO** circumstances will changes be allowed to any samples. **DO NOT** use addressograph labels on samples as the analysers are not compatible with these labels and they cannot be processed.

6.2.1. Blood Transfusion Sample Labelling Requirements

Sample identification is of critical importance in blood transfusion. The process of ordering blood for possible transfusion involves both a request for a laboratory investigation and also for a prescribed therapeutic product.

The vast majority of major transfusion complications (although rare) are caused by clerical errors and it is therefore important to follow procedures for patient identification.

It is the responsibility of the staff undertaking sample collection to ensure that the sample bottle is labelled accurately and signed and that the form is completed with all relevant information and is transported to the laboratory in a timely and correct manner.

All samples and requests for Blood Transfusion must have the following minimum identifiers:

- Surname
- Forename
- CHI or TJ number
- Date of birth

- Gender
- Signature of person taking the sample
- Date and time of sample

The Department will **NOT** process specimens that are incorrectly or inadequately labelled. In this event, the requesting location will be contacted by telephone, and a replacement specimen requested.

Sample bottles should **NEVER** be pre-labelled or completed away from the patient's side.

Any form of printed label **MUST NOT** be used on blood transfusion sample bottles. All blood transfusion sample bottles **MUST** be hand written and **MUST** be signed by the individual taking the sample.

6.2.2. Consent

Consent for the performing of an assay is implied when the patient agrees to have a sample taken for this purpose. Those assays which require specific consent (as required under legislation), this will be discussed by a clinical member of staff during the clinical appointment/visit.

6.3. Transportation of Samples

It is the responsibility of the staff undertaking sample collection to ensure that the sample is transported to the laboratory in a timely and correct manner.

6.3.1. Portering Services

Specimens are uplifted from the various clinical units by Porters on a regular basis throughout the day. To have samples collected urgently please contact the porters.

6.3.2. Vacuum Tube Specimen Delivery System

Vacuum tube systems are available for the transportation of laboratory specimens at the GGH, GRI and Stobhill sites. Please note that samples for platelet aggregation must not be sent via the tube system.

6.3.3. Primary Care Specimen Collection Service

This service is co-ordinated and managed by the Facilities Managers (**NOT** by the Department of Haematology), a specimen collection service operates, up to twice a day, for the routine collection and delivery of laboratory specimens between service users in General Practice and Primary Care Health Centres to the laboratories of the Department. All enquiries relating to these services should be directed to the relevant Facilities Manager.

6.3.4. Sending Specimens by Post

The Royal Mail supplies prepaid, single-use mailing containers (Safebox) that meets current legislation (UN3373) for posting laboratory specimens. Regardless of container type the following requirements apply for posted specimens:

- The primary container (specimen bottle) must be leak-proof and must not contain more than 500ml.

- There must be absorbent material, which must be present in sufficient quantity to absorb the entire content of the primary container, placed between the primary container and a secondary container.
- The secondary container must be leak-proof.
- The secondary packaging must not contain more than 4 litres (includes multiple primary containers placed into a single secondary container).
- Secondary container must be labelled with “Biological Material”, “Biohazardous Sample”, or similar, and must have the laboratory destination and return address clearly marked.

6.4. Restricted Specimens

Patients from whom specimens **MUST NOT BE SENT** without approval of an Infectious Disease/Control Clinician:

- Specimens from patients known or suspected to have SARS.
- Specimens from patients with possible or confirmed Viral Haemorrhagic Fever.
- Any other hazard category 4 pathogens

7. Assay Repertoire and Turn Around Times

The different haematological tests and bottle types (by colour) required are listed below: All turn-around times are for routine requests unless stated and are timed from receipt of the sample in the Laboratory as the collection and transportation of samples is out with the laboratories authority and control. Assays can be processed as urgent but the laboratory must be informed in advance. Assays out with our ISO15189 scope are indicated with two asterisks.

7.1. Haematology

Assay	Bottle Required	Site Performed at	Turnaround Time		
			Urgent*	In Patients	Outpatient/GPs
Full Blood Count (FBC)	4ml Purple	All	1hr	2hrs	4hrs
ESR	4ml Purple	GRI, GGH	1hr	2hrs	4hrs
Reticulocyte Count	4ml Purple	All	1hr	2hrs	4hrs
			Turnaround Time (All areas)		
Blood film	4ml Purple	GRI,GGH	24 hrs.		
Bone Marrow	4ml Purple	GRI,GGH	48 hrs.		
Body Fluid Analysis**†	Fluid dependent	GRI	4hrs		
Iron Stain**	4ml Purple	GRI,GGH	48 hrs.		
Glandular Fever Screen	4ml Purple	GGH, GRI	4 hrs.		
Malarial Parasites	4ml Purple	GGH, GRI	4 hrs.		
Sickle Cell Screen	4ml Purple	GGH, GRI	2 hrs.		
Kleihauer	4ml Purple	GRI	24 hrs.		
Haemoglobinopathy screen	4ml Purple	GRI	72 hrs.		
Urinary Haemosiderin**	20ml urine in sterile container	GRI, GGH	48 hrs.		

* Samples from A&E are considered Urgent all other areas are processed as routine unless contacted to process as urgent

** Assay not currently within our ISO15189 accredited scope

†Ascetic Fluid and CSF only

7.1.1. Malaria Parasite Screening Request Requirements

Laboratory investigation of malaria requires a full blood count (purple tube). It is essential that the

laboratory receives the sample the same day as it is taken.

A full travel history (country/countries recently visited) **MUST** be stated on the request. This can and will aid species identification.

Two investigations are performed – a rapid Malaria antigen screening test and thick and thin blood film examination for parasites. The rapid diagnostic test detects the presence of malaria only it **CANNOT** provide confirmation of species or level of parasitaemia. This can only be provided after examination of the blood films. Due to the limitations of the technology the Malaria RDT is unable to detect *P Knowlesi*.

In all cases where malaria is confirmed, the result will be telephoned to the requesting location by a member of the Haematology medical staff.

Malaria is a parasitic disease found in tropical and subtropical regions. It is caused by protozoa of the genus Plasmodium. Five species of this protozoa cause malaria in humans, *P.vivax*, *P.ovale*, *P.malariae*, *P Knowlesi* and *P.falciparum*. Of these infections *P.vivax*, *P.ovale* and *P.malariae* can cause severe illness, but *P.falciparum* and *P Knowlesi* can cause a much more serious illness which can be fatal. *P.falciparum* and *P Knowlesi* infections must be identified urgently. A full travel history of all countries and regions is therefore essential.

Malaria must be diagnosed without delay in order to commence appropriate treatment.

7.1.2. Haemoglobinopathy Screening Request Requirements

A Family Origin Questionnaire **MUST** accompany all requests for Haemoglobinopathy screening from ante-natal patients. These are not supplied by the laboratory but are generated using the PNBS system.

7.2. Haemostasis

Assay	Bottle Required	Site Performed at	Turnaround Time		
			Urgent*	In Patients	Outpatient/GPs
Prothrombin time	3.5ml Blue	GRI, GGH, Stobhill	1hr	2hrs	4hrs
APTT	3.5ml Blue	GRI, GGH, Stobhill	1hr	2hrs	4hrs
TCT	3.5ml Blue	GRI, GGH, Stobhill	1hr	2hrs	4hrs
Fibrinogen	3.5ml Blue	GRI, GGH, Stobhill	1hr	2hrs	4hrs
D-dimer	3.5ml Blue	GGH,GRI	1hr	2hrs	4hrs
INR	3.5ml Blue	GRI, GGH, Stobhill	1hr	2hrs	4hrs
APTT ratio	3.5ml Blue	GRI, GGH, Stobhill	1hr	2hrs	4hrs
PT, APTT, TCT using Mechanical clot detection method**	3.5ml Blue	GRI only	1hr	2hrs	4hrs
Protamine Sulphate (TCT)**	3.5ml Blue	GRI, GGH	1hr	2hrs	4hrs
Assay	Bottle Required	Site Performed at	Turnaround Time all Areas		
ADAMTS13 Activity	2 x 3.5 mL Blue	GRI	24hrs		
Anti Xa LMWH	3.5ml Blue	GRI	24 hrs		
Anti Xa UFH	3.5ml Blue	GRI	24 hrs		
Anti Xa Orgaran/danapariod	3.5ml Blue	GRI	24 hrs		
Anti Xa Fondaparinux	3.5ml Blue	GRI	24 hrs		
Apixaban level	3.5ml Blue	GRI	24 hrs		
Rivaroxaban level	3.5ml Blue	GRI	24 hrs		
Edoxaban level	3.5ml Blue	GRI	24 hrs		
Dabigatran level	3.5ml Blue	GRI	24 hrs		
Argatroban level	3.5ml Blue	GRI	24 hrs		

Assay	Bottle Required	Site Performed at	Turnaround Time all Areas
Heparin Induced Thrombocytopenia (HIT)	3.5ml Blue	GRI	By arrangement with consultant haematologist
Reptilase**	3.5ml Blue	GRI	By Arrangement***
Factor II	3.5ml Blue	GRI	14 days
Factor V	3.5ml Blue	GRI	14 days
Factor VII	3.5ml Blue	GRI	14 days
Factor VIII	3.5ml Blue	GRI	7 days
Factor IX	3.5ml Blue	GRI	7 days
Factor IX (Refixia)	3.5ml Blue	GRI	7 days
Factor X	3.5ml Blue	GRI	14 days
Factor XI	3.5ml Blue	GRI	7 days
Factor XII	3.5ml Blue	GRI	7 days
Factor XIII	3.5ml Blue	GRI	14 days
Fibrinogen Antigen**	3.5ml Blue	GRI	14 days
Chromogenic FVIII	3.5ml Blue	GRI	10 days
Emicizumab concentration**	3.5ml Blue	GRI	7 days
VWF:Ag	3.5ml Blue	GRI	10 days
VWF:RCo	3.5ml Blue	GRI	10 days
VWF:CBA	3.5ml Blue	GRI	10 days
Anti Cardiolipin Antibodies†	3.5ml Blue	GRI	14 days
Antithrombin activity	3.5ml Blue	GRI	7 days
Antithrombin Ag**	3.5ml Blue	GRI	28 days
Protein C activity	3.5ml Blue	GRI	7 days
Protein S (Free) Ag	3.5ml Blue	GRI	7 days
Platelet aggregation studies**	Contact Lab	GRI	By Arrangement***
Plasminogen	3.5ml Blue	GRI	42 days
α ₂ Antiplasmin	3.5ml Blue	GRI	42 days
Factor V Leiden	3.5ml Blue	GRI	28 days
ProthrombinG20210A Mutation	3.5ml Blue	GRI	14 days
CompleteThrombophilia Screen	4 x 3.5ml Blue	GRI	14 days
Inherited Thrombophilia Screen	2 x 3.5ml Blue	GRI	14 days
Acquired Thrombophilia Screen	2 x 3.5ml Blue	GRI	14 days
Investigation of Prolonged APTT	2 x 3.5ml Blue	GRI	7 days
Antiphospholipid Screen†	2 x 3.5ml Blue	GRI	14 days
Inhibitor to Factor VIII (Human)	3.5ml Blue	GRI	7 days
Inhibitor to Factor VIII (Porcine)	3.5ml Blue	GRI	7 days
Inhibitor to Factor VIII (Chromogenic)	3.5ml Blue	GRI	7 days
Inhibitor to FV	3.5ml Blue	GRI	7 days
Inhibitor to FIX	3.5ml Blue	GRI	7 days

* Samples from A&E are considered Urgent all other areas are processed as routine unless contacted to process as urgent

** Assay not currently within our ISO15189 accredited scope

*** By arrangement, tests only performed after approval by Dr R.Rodgers or Dr C. Bagot. These tests are performed infrequently and therefore have no defined Turnaround Time but will take significantly longer than other available assays.

† This assay has not been validated for paediatric samples and is not currently within our ISO15189 accredited scope for those samples.

7.2.1. Haemostasis Assays General Requirements

All samples **MUST** be received < 4 hours from time of venepuncture. All sample bottles must be filled correctly.

Samples for Platelet Aggregation assays **MUST** be pre-arranged with the laboratory.

Urgent requests will be processed within 24 - 48 hrs if clinically indicated and on agreement by a

Consultant Haematologist.

Please note that the results of Haemostasis tests may be affected by extremely high levels of haemoglobin, bilirubin, triglycerides, heparin, or rheumatoid factor. Levels are test specific – please contact the laboratory for further guidance.

7.2.2. Anticoagulation Therapy

The department offers a service for monitoring anticoagulation therapy which includes patients on Warfarin, Unfractionated Heparin, Low Molecular Weight Heparin, Fondaparinux, Danaparoid and Direct Oral Anticoagulants (DOACs). Clinical support and advice is available.

7.2.3. Anti Xa Assays Special Requirements

Requests for AXa UFH must be received in the laboratory within 1 hour of venepuncture

Patients on Low Molecular Weight Heparin must have the sample taken 3.5 to 4 hrs post dose. The type of Heparin must be stated on the request.

Patients on Direct Oral Anticoagulants (DOACs) must have the sample taken 3hrs post dose. The type of DOAC must be stated on the request.

7.2.4. Lupus Anticoagulant Assays Special Requirements

Samples must be received within 4 hours of the time of venepuncture.

7.2.5. Anticoagulation Service

Anticoagulant clinics, for monitoring of warfarin therapy, are run by the Glasgow and Clyde Anticoagulation Service (GCAS). A Clinical Nurse Specialist led community based service.

7.3. Haemato-oncology

The laboratory at Gartnavel General Hospital in association with the West of Scotland Regional Stem Cell Processing Service provides a comprehensive clinical Haemato-Oncology service which includes immunophenotyping.

Clinical support and advice is available.

Assay	Bottle Required	Site Performed at	Turnaround Time
CD34	4ml Purple EDTA	GGH	2 hrs
Immunophenotyping	4ml Purple EDTA	GGH	72 hrs
PNH	4ml Purple EDTA	GGH	24 hrs
Hereditary Spherocytosis (HS)**	4ml Purple EDTA	GGH	72 hrs*
Platelet Membrane Glycoproteins (PMG)**	3.5ml Blue Citrate	GGH	72 hrs*

*Prior arrangement necessary - Please phone 57707/57708 to discuss requirements.

**Assay not currently within our ISO15189 accredited scope

7.3.1. Haemato-oncology sampling requirements

All peripheral blood and bone marrow samples being sent for immunophenotyping should be labelled as detailed in section 6.2. All relevant clinical information and the timing of the sample must be

completed on the request form accompanying the sample. The laboratory must be made aware of all urgent samples that are being sent so they can be prioritised appropriately. Users will be notified of inappropriate requests or of samples not meeting acceptance criteria.

Samples must be sent with two representative smears which have been allowed to air dry. Samples must be transported to the laboratory as soon as possible and kept at ambient temperature.

Specimens requiring assay for platelet membrane glycoproteins or hereditary spherocytosis must be discussed with the lab prior to sampling and must be accompanied with the following controls.

Samples for Hereditary Spherocytosis must include six age matched control samples (with normal red cell indices) to be sent with HS Assay for individuals under 16 years of age.

Two control samples in blue citrate bottles are required with to be sent with a sample for the PMG Assay.

Bodily fluids and CSF can be sent in Universal Containers (2 to 3 ml is sufficient for analysis of cell markers). Anticoagulant is not necessary for these samples. CSF samples must be transported to the laboratory immediately for analysis.

7.4. Blood Transfusion

Blood Transfusion Maximum Surgical Blood Ordering Schedule (MSBOS), policies, guidelines and forms, both local and pan NHS GG&C, are available on the Blood Transfusion page of the Intranet.

Assay	Bottle Required	Site Performed at	Turnaround Time
Group and Antibody Screen	6ml Pink	GRI /GGH	4 hrs
Cross Match	6ml Pink	GRI/GGH	2 hrs
Antibody Identification	6ml Pink	GRI/GGH	4 hrs

7.4.1. Routine Blood Product Orders

Blood requirements for surgery vary depending on the procedure. For procedures likely to require blood products a pre-determined number of units should be requested accordingly to the MSBOS and should be requested in good time prior to surgery.

In those cases where blood is rarely required a group and screen is recommended. A second confirmatory sample will be required to be processed before the issue of any blood products if the patient has not previously been grouped.

The Transfusion samples are stored in the laboratory for 7 days, unless they are a preoperative sample with a given date for the operation in which case the samples will be stored in the laboratory for 14 days. Samples will be stored for 72 hours only, if the patient is pregnant or has been transfused in the past three months. If blood is subsequently required it can be provided following compatibility testing or Electronic Issue. The Blood Transfusion Laboratory must be informed and a crossmatch request form must be supplied to the laboratory to confirm the request for blood products.

Blood Transfusion MSBOS, policies, guidelines and forms both local and pan NHS GG&C are available on the Blood Transfusion page of the Intranet.

7.4.2. Emergency Cross Match Requests

The Blood transfusion laboratory should be telephoned in advance, and the sample transported to the laboratory immediately and not left to the routine specimen collection. It will be useful if the degree of emergency is stated. Out with laboratory core hours the on duty BMS **MUST** be contacted

7.4.3. Atypical Red Cell Antibodies

Occasionally patients may have antibodies to blood group antigen systems other than ABO and Rhesus. These may have been identified on a previous occasion, in which case the patient may have been issued with an antibody card indicating the identity of the antibody/antibodies.

It may be more difficult to provide compatible blood for these patients and requirements should be discussed with the Blood Transfusion Laboratory. As much advanced planning and notice as is possible of operations for such patients should be undertaken.

Failure to do so will lead to a delay in the provision of blood products.

In situations where further blood is required for a patient who has already been recently transfused a fresh sample must be sent for cross matching.

7.4.4. Transfusion Reactions

Mild transfusion reactions are not uncommon, however severe life-threatening haemolytic reactions are rare.

Transfusion reactions can be classified as:

- Haemolytic Reactions
- Non-haemolytic febrile transfusion reactions
- Urticarial and Anaphylactic reactions

Should a transfusion reaction be suspected immediately contact a Haematology Registrar or Consultant. A Transfusion reaction form must be completed and these are available on the departmental intranet page and webpage on the NHSGGC website.

7.4.5. Blood Components

The following Blood components can be requested, some must be discussed with a Haematology clinician prior to requesting.

7.4.5.1. Red Cell Concentrate

Packs contain approximately 200mls of concentrated red cells from plasma reduced donor units or plasma depleted units. All red cell products in the UK are leucodepleted.

7.4.5.2. Fresh Frozen Plasma

This is normally issued after discussion with a Haematology clinician and must be given immediately on delivery to the ward/unit. First dose FFP in the context of correction of Haemostasis (for patient's not on warfarin) can be requested without contacting a haematology clinician, provided the cause of laboratory abnormalities is understood.

7.4.5.3. Platelet Concentrates

This is normally issued after discussion with a Haematology clinician, there may be some delay in receiving the product as normally it is necessary to order and have it delivered from SNBTS. It **MUST NOT** be refrigerated.

Cryoprecipitate

This is normally issued after discussion with a haematology clinician and should be given immediately on delivery to the ward/unit.

7.4.5.4. Anti-D Immunoglobulin

Available in various doses and issued dependent on the stage of pregnancy or in the presence of, or possibility of, Foetal Maternal Haemorrhage (FMH).

7.4.5.5. Specific Haemostasis factor concentrates

These are issued after discussion with a haematology clinician.

7.5. Costs

Information about the costs and charges of our assay repertoire can be obtained from the Technical Services Manager on application. Their contact details can be found in section 4

8. Referred Assays

The following assays are referred to laboratories within or out with NHS GG&C. All laboratories are accredited to ISO15189 unless stated.

Assay	Referral Laboratory Location	Turnaround Time
Malaria Screen Confirmation	Microbiology Glasgow Royal Infirmary	48hrs
Haemoglobinopathy Confirmation	Oxford Molecular Diagnostic Laboratory	28 days
Haemoglobinopathy Confirmation	Western General Hospital Edinburgh	28 days
EPO receptor and VHL genetic Analysis	Belfast City Hospital, Belfast	28 days
FIP1L1-PDGFRa c-KIT d816v mutation	Salisbury District Hospital, Salisbury	28 days
Chromosome Breakage	Guy's and St Thomas' Hospital, London	10 days
Fibrinogen Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Factor V Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Factor VII Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Factor VIII Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Factor IX Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Factor X Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Factor XI Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Factor V & VIII combined Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Antithrombin Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Protein C Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Protein S Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
VWF Multimers	Royal infirmary, Edinburgh	84 days
VWF Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
ADAMST13 Antibodies	HSL, London	28 days
Platelet Nucleotides	Royal infirmary, Edinburgh	28 days
Platelet (MYH9) Genetic Analysis	Royal infirmary, Edinburgh	42-56 days
Glanzmann's Thrombasthaenia Genetics	Royal infirmary, Edinburgh	42-56 days
Bernard Soulier Syndrome Genetics	Royal infirmary, Edinburgh	42-56 days
X-Matching (multiple red cell antibodies)	SNBTS Gartnavel	4 hrs**

Assay	Referral Laboratory Location	Turnaround Time
Red Cell Antibody Identification (multiple red cell antibodies)	SNBTS Gartnavel	7 days
HLA Screening (Platelets and WBC)	SNBTS Gartnavel	48 hrs
Anti IgA antibodies	SNBTS Gartnavel	48 hrs
Neutrophil and Granulocyte Antibodies	NHSBT Bristol	48 hrs
Anti D Quantification	SNBTS Gartnavel	4 days
Foetal Maternal Haemorrhage quantification	SNBTS Gartnavel	48 hrs
Foetal Genotyping	NHSBT Bristol	10 days
Transfusion Reaction Investigations (Requested by Haematologist)	SNBTS Gartnavel	7-14 days

* Performed only by request no available turnaround time due to assay frequency.

** For very rare antibody types/combinations this may be considerably longer especially if red cell antibody identification is required as well.

9. Reference Ranges

The Reference Ranges are for guidance only, and are derived from *Dacie and Lewis Practical Haematology - 12th Edition* unless stated. Advice, both clinical and technical is available by contacting either a member of the Haematology clinical or technical staff respectively. Some ranges differ due to age or sex. Some assays do not have a numerical reference range with the result of the assay being given in a clinical comment or as a statement of positivity or negativity. Others have a therapeutic range. For those tests that have a therapeutic range please consult the appropriate policy or contact a member of the Haematology clinical team if guidance is required. All ranges are for adults unless stated. All ranges are for both males and females unless stated.

9.1. Haematology Reference Ranges

The Adult, Infant and Children's Reference Ranges are sourced from, *Dacie and Lewis Practical Haematology 12th Ed (2017)*: S M Lewis, B J Bain, I Bates, M Laffan.

The Infant and Children's White blood cell and differential Reference Ranges are sourced from *Pediatric Hematology 3rd Ed (2006)*: R J Arceci, I M Hann, O P Smith.

Unless stated otherwise.

9.1.1. Haematology Reference Ranges for Adults

Assay	Male Reference Range	Female Reference range
WBC	4.00 – 10.00 x 10 ⁹ /L	
Haemoglobin*	130 - 180g/L	115 - 165g/L
RBC*	4.50 – 6.50 x 10 ¹² /L	3.80 - 5.80 x 10 ¹² /L
HCT*	0.400 - 0.540	0.370 - 0.470
MCV	83 - 101fl	
MCH	27 - 32pg	
MCHC	315 -345 g/L	
Platelets	150 - 410 x 10 ⁹ /L	
Neutrophils	2.00 - 7.00 x 10 ⁹ /L	
Lymphocytes*	1.1 – 5.0 x 10 ⁹ /L	
Monocytes	0.20 – 1.0 x 10 ⁹ /L	
Eosinophils	0.02 – 0.5 x 10 ⁹ /L	
Basophils	0.02 – 0.10 x 10 ⁹ /L	
Reticulocytes	50 – 100 x 10 ⁹ /L	
Blood Film	Comment	
Glandular Fever Screen	Comment	
Malaria Parasite Screen	Comment	
Urinary Haemosiderin	Comment	

*Reference range by local expert review

9.1.1. Haematology Reference Ranges for Children

Assay	1 Year	2 to 6 Years	6 to 12 Years	12 to 18 Years
WBC	6.00 – 16.00 x 10 ⁹ /L	6.00 – 17.00 x 10 ⁹ /L	4.50 – 13.00 x 10 ⁹ /L	4.5 – 13.00 x 10 ⁹ /L
Haemoglobin	111 - 141g/L	110 - 140g/L	115 - 155g/L	115 - 155g/L
RBC	3.90 – 5.10 x 10 ¹² /L	4.00 – 5.20 x 10 ¹² /L	4.00 – 5.20 x 10 ¹² /L	4.00 – 5.20 x 10 ¹² /L
HCT	0.300 - 0.380	0.340 - 0.400	0.350 - 0.450	0.350 - 0.450
MCV	72 - 84fL	75 - 87fL	77 - 95fL	77 - 95fL
MCH	25 - 29pg	24 - 30pg	25 - 33pg	25 - 33pg
MCHC	320 -360 g/L	310 -370 g/L	310 -370 g/L	310 -370 g/L
Platelets	200 - 550 x 10 ⁹ /L	200 - 490 x 10 ⁹ /L	170 - 450 x 10 ⁹ /L	170 - 450 x 10 ⁹ /L
Neutrophils	1.00 - 8.00 x 10 ⁹ /L	1.50 - 8.50 x 10 ⁹ /L	1.50 - 8.00 x 10 ⁹ /L	1.50 - 6.00 x 10 ⁹ /L
Lymphocytes	3.40 - 10.5 x 10 ⁹ /L	1.80 - 8.40 x 10 ⁹ /L	1.50 - 5.0 x 10 ⁹ /L	1.0 – 4.50 x 10 ⁹ /L
Monocytes	0.20 – 0.9 x 10 ⁹ /L	0.15 – 1.30 x 10 ⁹ /L	0.15 – 1.30 x 10 ⁹ /L	0.15 – 1.30 x 10 ⁹ /L
Eosinophils	0.05 – 0.9 x 10 ⁹ /L	0.05 – 1.10 x 10 ⁹ /L	0.05 – 1.0 x 10 ⁹ /L	0.05 – 0.80 x 10 ⁹ /L
Basophils	0.02 - 0.13 x 10 ⁹ /L	0.02 - 0.12 x 10 ⁹ /L	0.02 - 0.12 x 10 ⁹ /L	0.02 - 0.12 x 10 ⁹ /L
Reticulocytes	30 - 100 x 10 ⁹ /l	30 - 100 x 10 ⁹ /L	30 - 100 x 10 ⁹ /L	30 - 100 x 10 ⁹ /L
Blood Film	Comment	Comment	Comment	Comment
Glandular Fever Screen	Comment	Comment	Comment	Comment
Malaria Parasite Screen	Comment	Comment	Comment	Comment

9.1.2. Haematology Reference Ranges for Infants

Assay	Birth	Day 3	Day 7
WBC	10.00 – 26.00 x 10 ⁹ /L	10.00 – 26.00 x 10 ⁹ /L	10.00 – 26.00 x 10 ⁹ /L
Haemoglobin	140 - 220g/L	150 - 210g/L	135 - 215g/L
RBC	5.00 – 7.00 x 10 ¹² /L	4.00 – 6.60 x 10 ¹² /L	3.90 – 6.30 x 10 ¹² /L
HCT	0.450 - 0.750	0.450 - 0.670	0.420 - 0.660
MCV	100 - 120fL	92 - 118fL	88 - 126fL
MCH	31 - 37pg	31 - 37pg	31 - 37pg
MCHC	300 -360 g/L	290 -370 g/L	280 -380 g/L
Platelets	100 - 450 x 10 ⁹ /L	210 - 500 x 10 ⁹ /L	160 - 500 x 10 ⁹ /L
Neutrophils	2.70 - 14.40 x 10 ⁹ /L	2.70 - 14.40 x 10 ⁹ /L	2.70 - 14.40 x 10 ⁹ /L
Lymphocytes	2.0 – 7.3 x 10 ⁹ /L	2.0 – 7.3 x 10 ⁹ /L	2.0 – 7.3 x 10 ⁹ /L
Monocytes	0.0 – 1.9 x 10 ⁹ /L	0.0 – 1.9 x 10 ⁹ /L	0.0 – 1.9 x 10 ⁹ /L
Eosinophils	0.0 – 0.85 x 10 ⁹ /L	0.0 – 0.85 x 10 ⁹ /L	0.0 – 0.85 x 10 ⁹ /L
Basophils	0.0 - 0.10 x 10 ⁹ /L	0.0 - 0.10 x 10 ⁹ /L	0.0 - 0.10 x 10 ⁹ /L
Reticulocytes	120 - 400 x 10 ⁹ /L	50 - 350 x 10 ⁹ /L	50 - 100 x 10 ⁹ /L
Blood Film	Comment	Comment	Comment
Malaria Parasite Screen	Comment	Comment	Comment

Assay	Day 14	1 Month	2 Months	6 Months
WBC	6.00 – 21.00 x 10 ⁹ /L	6.00 – 21.00 x 10 ⁹ /L	5.00 – 15.00 x 10 ⁹ /L	6.00 – 17.00 x 10 ⁹ /L
Haemoglobin	125 - 205g/L	115 - 165g/L	94 - 130g/L	111 - 141g/L
RBC	3.90 – 6.20 x 10 ¹² /L	3.00 – 5.40 x 10 ¹² /L	3.10 – 4.30 x 10 ¹² /L	4.10 – 5.30 x 10 ¹² /L
HCT	0.310 - 0.710	0.330 - 0.530	0.280 - 0.420	0.300 - 0.400
MCV	86 - 124fL	92 - 116fL	87 - 103fL	68 - 84fL
MCH	31 - 37pg	30 - 36pg	27 - 33pg	24 - 30pg
MCHC	280 -380 g/L	290 -370 g/L	285 -355 g/L	300 -360 g/L
Platelets	170 - 550 x 10 ⁹ /L	210 - 500 x 10 ⁹ /L	210 - 650 x 10 ⁹ /L	200 - 550 x 10 ⁹ /L
Neutrophils	1.50 - 5.40 x 10 ⁹ /L	1.50 - 5.40 x 10 ⁹ /L	0.70 – 4.80 x 10 ⁹ /L	1.00 - 6.00 x 10 ⁹ /L
Lymphocytes	2.80 - 9.10 x 10 ⁹ /L	2.80 - 9.10 x 10 ⁹ /L	3.3 – 10.3 x 10 ⁹ /L	3.30 – 11.5 x 10 ⁹ /L
Monocytes	0.10 – 1.7 x 10 ⁹ /L	0.10 – 1.7 x 10 ⁹ /L	0.40 – 1.2 x 10 ⁹ /L	0.20 – 1.3 x 10 ⁹ /L
Eosinophils	0.0 – 0.85 x 10 ⁹ /L	0.0 – 0.85 x 10 ⁹ /L	0.05 – 0.90 x 10 ⁹ /L	0.1 – 1.10 x 10 ⁹ /L
Basophils	0.0 - 0.10 x 10 ⁹ /L	0.0 - 0.10 x 10 ⁹ /L	0.02 - 0.13 x 10 ⁹ /L	0.02 - 0.13 x 10 ⁹ /L
Reticulocytes	50 - 100 x 10 ⁹ /L	20 - 60 x 10 ⁹ /L	30 - 50 x 10 ⁹ /L	40 - 100 x 10 ⁹ /L
Blood Film	Comment	Comment	Comment	Comment
Malaria Parasite Screen	Comment	Comment	Comment	Comment

9.1.3. ESR

Test	17-50 Years	50-61 Years	61-70 Years	>70 Years
ESR (male) mm/hr	≤ 10	≤ 12	≤ 14	≤ 30
ESR (female) mm/hr	≤ 12	≤ 19	≤ 20	≤ 35

9.2. Haemoglobinopathy Assays

The Reference Ranges are sourced from Haemoglobinopathy Diagnosis 2nd Ed: B J Bain

Assay	Adult Reference Range
HbF	<1.0%
HbA2	2.0 – 3.5%
Sickle Cell RDT	Comment

9.3. Haemato-Oncology

9.3.1. Immunophenotyping

Immunophenotyping reference ranges are considered not to be appropriate and an interpretative comment is provided on the report which provides all the relevant clinical, morphological and immunophenotyping data.

CD34 absolute values are evaluated by the consultant in charge of apheresis with particular regard to the timing of mobilisation and therefore a specific reference range for CD34 counts is not applicable either.

9.4. Haemostasis Reference Ranges

All Haemostasis reference ranges are locally derived from a pool of normal individuals.

9.4.1. Routine Haemostasis Ranges

Assay	Adult Reference Range
Prothrombin time	9 – 13 sec
APTT	27 – 36 sec
TCT	11 – 15 sec
Fibrinogen	1.7 – 4.0 g/L
D-Dimer	<243ng/mL
D-Dimer for exclusion of VTE	<230ng/mL
INR	Therapeutic range (2.0 -4.0 Depending on Indication)
APTT ratio	Therapeutic range 1.8 – 2.8
Anti Xa.	Therapeutic ranges for individual anticoagulants

9.4.2. Specific Haemostasis Assay Reference Ranges

Assay	Male Reference Range	Female Reference Range
Reptilase	13 – 20 sec	
Factor II	97 - 141 iu/dL	
Factor V	66 - 167 iu/dL	
Factor VII	67 - 153 iu/dL	
Factor VIII	58 - 152 iu/dL	
Factor IX	81 - 157 iu/dL	
Factor X	79 - 155 iu/dL	
Factor XI	82 - 151 iu/dL	
Factor XII	59 - 164 iu/dL	
Factor XIII	70 - 140 iu/dL	
Fibrinogen Antigen	1.8 – 3.4 g/dL	
Chromogenic FVIII	50 – 200 iu/dL	
VWF:Ag	51 - 170 iu/dL	
VWF:RCo	46 – 166 iu/dL	
VWF:CBA	50-160 iu/dL	
DRVVT Screen	0.87 – 1.21 ratio	
DRVVT Confirm	0.81 – 1.08 ratio	
ACL Antibody (IgG)*	<20.0 U/mL	
ACL Antibody (IgM)*	<20.0 U/mL	
Antithrombin activity	82 - 123 iu/dL	
Antithrombin Ag	75 – 128 iu/dL	
Assay	Male Reference Range	Female Reference Range
Protein C activity	71 - 146 iu/dL	
Protein S (Free) Ag	75 - 148 iu/dL	65 - 137 iu/dL
Platelet aggregation	Comment	
Plasminogen	80 – 133 U/dL	
α ₂ Antiplasmin	98 – 122 U/dL	
Factor V Leiden	Comment	
Prothrombin G20210A Mutation	Comment	
Factor VIII Inhibitor Assay	Comment	
ADAMTS13 Activity	60.6 – 130.6 IU/dL	

*Please note that references ranges for ACL Antibody (IgM/G) are not validated for paediatric samples.

9.5. Haematinics Assays

Haematinic assays are processed by the Biochemistry department but are reported within the Haematology section of SCI store, IS and Clinical Portal. Clinical advice is available from the Haematology Clinical staff detailed in section 4.3.6. Technical advice is available from the biochemistry department.

Assay	Male	Female
B12 (active)	≥25 pmol/L	
B12 (active) Indeterminate Level	25– 70 pmol/L	
Serum Folate	3.1 – 20 ug/L	
Ferritin	15 – 300 ug/L	15 – 200 ug/L
EPO	2.6-18.5 U/L	

9.6. Uncertainty of Measurement (UoM)

No measurement is exact. When something is measured, the outcome depends on the measuring system, the measurement procedure, the skill of the operator, the environment, and other effects. Even if the item were to be measured several times, in the same way and in the same circumstances, a different measured value would, in general be obtained each time, assuming the measuring system has sufficient resolution to distinguish between the values. This variability, for those results that are expressed numerically, has been calculated and is given in the column UoM. These values have been reviewed by the clinical staff and have been deemed not to be sufficient to affect any clinical decisions that may be taken using the results of analysis.

The uncertainty of measurement is expressed as a 95% confidence interval unless stated.

The range stated in the tables is **NOT** the assay reference range but the high and low values between which the uncertainty of measurement estimation has been established.

All uncertainty of Measurement estimations have been reviewed by a consultant clinical staff member and considered not to affect any decision required for patient care.

Uncertainty of Measurement must not be confused with error. These are defined as:

- **Uncertainty of measurement:** Quantified doubt about the result of a measurement
- **Error:** The difference between the measured value and the true value of the object being measured.

9.6.1. Haematology UoM

Assay	Uncertainty of Measurement +/-	Range	
		From	To
WBC	0.12 x10 ⁹ /L	2.65 x10 ⁹ /L	3.62 x10 ⁹ /L
	0.21 x10 ⁹ /L	6.34 x10 ⁹ /L	8.49 x10 ⁹ /L
	0.36 x10 ⁹ /L	14.94 x10 ⁹ /L	19.77 x10 ⁹ /L
Haemoglobin	1.1 g/L	55 g/L	64 g/L
	1.5 g/L	116 g/L	130 g/L
	1.9 g/L	152 g/L	172 g/L
RBC	0.04 x10 ¹² /L	2.19 x10 ¹² /L	2.48x10 ¹² /L
	0.06 x10 ¹² /L	4.13 x10 ¹² /L	4.68 x10 ¹² /L
	0.07 x10 ¹² /L	4.95 x10 ¹² /L	5.65 x10 ¹² /L
HCT	0.005	0.155	0.197
	0.009	0.319	0.406
	0.010	0.418	0.552
MCV	1.6 fL	70.9 fL	81.3 fL
	1.5 fL	77.1 fL	90.9 fL
	1.6 fL	82.4 fL	93.7 fL
MCH	0.5 pg	22.5 pg	28.8 pg
	0.4 pg	24.8 pg	31.2 pg
MCH	0.5 pg	27.6 pg	33.4 pg
MCHC	9.1 g/L	280 g/L	392 g/L
	7.3 g/L	291 g/L	391 g/L
	7.5 g/L	298 g/L	398 g/L
Platelets	5 x10 ⁹ /L	53 x10 ⁹ /L	125 x10 ⁹ /L
	11 x10 ⁹ /L	203 x10 ⁹ /L	291 x10 ⁹ /L
	18 x10 ⁹ /L	496 x10 ⁹ /L	635 x10 ⁹ /L
Neutrophils	0.06 x10 ⁹ /L	0.86 x10 ⁹ /L	1.39 x10 ⁹ /L
	0.14 x10 ⁹ /L	2.44 x10 ⁹ /L	3.45 x10 ⁹ /L
	0.31 x10 ⁹ /L	6.28 x10 ⁹ /L	8.83 x10 ⁹ /L
Lymphocytes	0.07 x10 ⁹ /L	0.55 x10 ⁹ /L	1.67 x10 ⁹ /L
	0.12 x10 ⁹ /L	1.56 x10 ⁹ /L	2.88 x10 ⁹ /L
	0.22 x10 ⁹ /L	2.86 x10 ⁹ /L	5.75 x10 ⁹ /L
Monocytes	0.07 x10 ⁹ /L	0.07 x10 ⁹ /L	0.76 x10 ⁹ /L
	0.09 x10 ⁹ /L	0.31 x10 ⁹ /L	1.50 x10 ⁹ /L
	0.19 x10 ⁹ /L	0.91 x10 ⁹ /L	3.12 x10 ⁹ /L
Eosinophils	0.04 x10 ⁹ /L	0.12 x10 ⁹ /L	0.44 x10 ⁹ /L
	0.09 x10 ⁹ /L	0.34 x10 ⁹ /L	1.10 x10 ⁹ /L
	0.24 x10 ⁹ /L	0.86 x10 ⁹ /L	2.81 x10 ⁹ /L
Basophils	0.01 x10 ⁹ /L	0.03 x10 ⁹ /L	0.28 x10 ⁹ /L
	0.03 x10 ⁹ /L	0.07 x10 ⁹ /L	0.73 x10 ⁹ /L
	0.08 x10 ⁹ /L	0.16 x10 ⁹ /L	1.83 x10 ⁹ /L
Reticulocytes	7.43 x10 ⁹ /L	78.9 x10 ⁹ /L	170.6 x10 ⁹ /L
	5.69 x10 ⁹ /L	57.2 x10 ⁹ /L	122.7 x10 ⁹ /L
Reticulocytes	3.25 x10 ⁹ /L	23.5 x10 ⁹ /L	59.3 x10 ⁹ /L
ESR	1 mm/hr	0 mm/hr	10 mm/hr
	1 mm/hr	50 mm/hr	60mm/hr

9.6.2. Manual Leucocyte Differential UoM

Assay	Uncertainty of Measurement +/-	Range	
		From	To
Neutrophils	6.59 %	48%	98%
Lymphocytes	6.14 %	1%	46%
Monocytes	4.36 %	1%	26%
Eosinophils	1.76 %	1%	9%
Basophils	0.75 %	0%	1%

9.6.3. Malaria Parasite Parasitaemia UoM

Assay	Uncertainty of Measurement +/-	Range	
		From	To
Malaria Parasitaemia	1.65%	0.3%	27.8%

9.6.4. Bone Marrow Myelogram

Cell Type	Uncertainty of Measurement +/-
Myeloblasts	1.7%
Proyelocytes	1.5%
Myelocytes	6.1%
Metamyelocytes	6.2%
Neutrophils	6.5%
Eosinophils	3.6%
Basophils	0.8%
Monocytes	3.6%
Erythrocytes	4.0%
Lymphocytes	5.3%
Plasma cells	1.2%

9.6.5. Haemoglobinopathy UoM

Assay	Uncertainty of Measurement +/-	Range	
		From	To
HbF	0.04 %	1.4 %	2.2 %
	0.10 %	8.6 %	9.6 %
HbA2	0.04 %	2.1 %	3.0 %
	0.07 %	5.2 %	6.0 %

9.6.6. Routine Haemostasis UoM

Assay	Uncertainty of Measurement +/-	Range	
		From	To
Prothrombin time	0.1 sec	9.2	13.2
	0.3 sec	18.4	27.6
APTT	0.4 sec	25.4	33.4
	0.4 sec	38.4	51.9
TCT	0.2 sec	11.7	17.7
	0.3 sec	15.9	21.6
Fibrinogen	0.1 g/L	1.6	2.6
	0.1 g/L	2.52	3.72
D-Dimer	9 ng/mL	204	340
	17 ng/mL	466	698
LMW Heparin	0.03U/mL	1.21	1.85
	0.02 U/mL	1.46	0.78
UF Heparin	0.01 U/mL	1.0	1.4
	0.005 U/mL	0.3	0.5
Apixaban	2.2 ng/mL	243	328
	1.7 ng/mL	55	85
Rivaroxaban	3.5 ng/mL	234	350
	1.7 ng/mL	61	102
Orgaran	0.02 U/mL	1.0	1.4
	0.01 U/mL	0.3	0.5
Fondaparinox	0.08 ug/mL	1.0	1.4
	0.04 ug/mL	0.3	0.5
Emicizumab	0.8 mg/mL	18.7	28.3
	2.0 ug/mL	55.8	83.8
Reptilase	0.7 sec	13.8	15.8
	0.7 sec	14.2	16.2

9.6.7. Special Haemostasis UoM

Assay	Uncertainty of Measurement +/-	Range	
		From	To
Factor II	1.2 iu/dL	77	117
	0.4 iu/dL	22	42
Factor V	2.2 iu/dL	79	119
	0.9 iu/dL	24	44
Factor VII	1.3 iu/dL	65	105
	0.4 iu/dL	19	39
Factor VIII	1.2 iu/dL	77	117
	0.6 iu/dL	16	36
Factor IX	2.8 iu/dL	83	123
	0.8 iu/dL	20	40
Factor X	1.6 iu/dL	70	110
	0.8 iu/dL	27	47
Factor XI	1.6 iu/dL	67	107
	0.5 iu/dL	21	41

Assay	Uncertainty of Measurement +/-	Range	
		From	To
Factor XII	3.1 U/dL	70	110
	2.6 iu/dL	23	43
Factor XIII	1.5 iu/dL	48	78
	1.0 iu/dL	13	33
Fibrinogen Antigen	0.1 g/L	1.82	3.39
Chromogenic FVIII	5.7 iu/dL	77	117
	1.3 iu/dL	16	36
VWF:Ag	1.4 iu/dL	100	130
	0.7 iu/dL	21	41
VWF:RCo	0.4 iu/dL	20	31
	1.6 iu/dL	75	117
VWF:CBA	0.8 iu/dL	19	30
	3.4 iu/dL	70	109
Lupus (APTT)	0.3 sec	28	33
DRVVTs	0.34 sec	30 sec	34 sec
DRVVTc	0.3 sec	30 sec	34 sec
ACL Antibody (IgG)	0.35 U/mL	6	12
	3.44 U/mL	69	120
ACL Antibody (IgM)	0.21 U/mL	6	11
	1.26 U/mL	52	120
Antithrombin activity	0.7 iu/dL	85	115
	0.6 iu/dL	16	36
Antithrombin Ag	2.9 iu/dL	85	115
	1.9 iu/dL	19	36
Protein C activity	0.6 iu/dL	86	116
	0.9 iu/dL	18	38
Protein S (Free) Ag	0.9 iu/dL	76	106
	0.8 iu/dL	19	39
APC Resistance +APC	2.9 sec	65	134
	0.4 sec	29	43
APC Resistance -APC	0.6 sec	30	50
	1.4 sec	50	91
Plasminogen	2.2 iu/dL	82	112
	1.6iu/dL	19	39
Antiplasmin	4.7iu/dL	85	115
	1.9	32	52
HIT	0.06	2.4	3.8
	0.03	0.4	0.7
ADAMTS13	2.2	24	46
	7.5	62	144

9.6.8. Haematinics Assays UoM

Haematinics uncertainty of measurement information has been supplied by Biochemistry GRI.

Assay	Uncertainty of Measurement +/-
B12	11.3 ng/L
Folate	1.47 ng/L
Folate	9.40 ng/L
Ferritin	0.31 ug/L
Ferritin	0.51 ug/L
EPO	5 U/L

9.6.9. Haemato-Oncology UoM

Assay	Uncertainty of Measurement +/-	Range	
		From	To
CD34	0.5 cells/uL	7 cells/uL	44 cells/uL

9.6.10. Qualitative Assays UoM

Uncertainty of measurement assessment for quantitative assays is assessed using a risk based method. Due to their composition and size they are not included in this manual. They are available on application to the quality manager.

The available assessments are:

- Factor V Leiden
- Prothrombin G20210A
- Acid Gel Haemoglobin Electrophoresis
- Cellulose Acetate Haemoglobin Electrophoresis
- Sickle Cell Solubility
- Blood Transfusion Qualitative tests
- Immunophenotyping
- Malaria RDT
- Glandular Fever RDT

9.7. Factors That Will Affect the Accuracy of Results

The following factors will affect the accuracy of the results however their total effect on the inaccuracy of the result cannot be quantified and in some cases will lead to an inability to perform the assay. Please note that some causes of variation are common to all sample types and requests and that they cannot be controlled by the laboratory but must be controlled by the individual service users or others. It is very important to identify and minimize significant pre-assay and post-assay conditions that will affect the accuracy of the assays.

The following are lists of conditions or situations that will affect results.

9.7.1. Common Factors

- Differences in patient preparation.
- Specimen collection technique.
- Transportation of sample.
- Storage time and storage temperature of sample within and out with the laboratory.
- Intra-individual variability (such as pregnancy, fasting/non-fasting, drug use, diurnal and underlying condition).
- Within individual biological variation.
- Environmental conditions within the laboratory.
 - Temperature, humidity and dust that may affect analysers, assays and sample stability.
- Reporting.
 - Number of significant figures

Further conditions will be more specific to a request as given below.

9.7.2. Blood Transfusion

- Age of sample - only valid for 24hrs at room temperature.
- Expiry date on sample.
- Haemolysed Samples.
- Lipaemic samples.
- Clotted samples.
- Insufficient samples.
- Recent Transfusion of blood of different group.
- Haemopoetic Stem cell transplant/BMT from donor of different blood group.

9.7.3. Haemostasis

- Haemostasis samples should preferably be taken before other test samples are drawn to avoid possible cross contamination of anticoagulants.
- Sodium citrate samples only must be used.
- Good venepuncture – poor venepuncture may lead to activation of the sample.
- No less than 90% fill
- No more than 110% fill.
- Blood must not be transferred from any other collection tube type to a Sodium citrate tube.
- Poor mixing of sample and anticoagulant to prevent clotting.
- High HCT's >0.55 will require an adjustment of sodium citrate volume – contact the laboratory for advice.
- Samples must be kept at room temperature and transported to the laboratory within 1hour of collection.
 - They must not be stored or transported on ice.
- Lipaemic samples.
- Icteric samples.
- Haemolysed samples.
- Residual Thrombin potential of factor deficient plasmas

9.7.4. Haematology

- 4ml EDTA samples only
- Age of sample - only valid for 24hrs
- Expiry date on sample.
- Haemolysed Samples.
- Lipaemic samples.
- Clotted samples.
- Insufficient samples.
- Haemoglobin level (sickle solubility test)
- Red blood cell count (ESR)
- Temperature (ESR)
- Due to limitations of the technology the Malaria RDT is unable to detect *P Knowlesi*.

9.7.5. Haemato-Oncology

- Clotted Samples
- Samples greater than 48 hours old.
- Samples for PNH analysis, must be received by the lab within 48hours of sampling as cells start to lose relevant CD markers
- Better quality bone marrow samples are achieved by drawing less than 3 ml of marrow as the volume of an aspirate is inversely proportional to its purity.
- Peripheral blood may haemodilute marrow aspirates and lead to differences in cell % and immunophenotype when flow cytometry is compared with morphology or histology.
- Bone marrow samples from patients with fibrosis may yield discrepant results,
- Deposition of reticulin or lysis of fragile cell populations in bone marrow samples may not lead to the detection of the presence of lymphoma by flowcytometry.
- CSF samples may contain few malignant cells amongst a reactive infiltrate and need to be received by the laboratory as soon as possible as cells deteriorate rapidly in this fluid.

10. Confidentiality

All data held within the department is done so in compliance with the NHSGGC Confidentiality and Data Protection Policy, the NHSGGC Confidentiality Policy, the NHS Code of Practice on Protecting Patient Confidentiality, The General Data Protection Regulations, The Data Protection Act 2018, The Blood Safety and Quality Regulations, The Medicines for Human Use (Clinical Trials) Regulations 2004.

11. Retention of Records

Records and Specimens are also held in compliance with the RCPATH guidelines on the Retention and Storage of Pathological Records and Specimens - 2015, 5thEd, The Data Protection Act 2018 and the requirements of The Medicines for Human Use (Clinical Trials) Regulations 2004 and the Human Tissues act 2004.

12. References

- The Medicines for Human Use (Clinical Trials) Regulations 2004
- Blood Safety and Quality Regulations 2005 and Amendment 2007
- Data Protection Act 2018

- Human Tissues Act 2004
- ISO 15189 – 2012: Medical Laboratories, Requirements for Quality & Competence
- Rules and Guidance for Pharmaceutical Manufacturers and Distributors: MHRA, 2017
- Good Clinical Practice Guide: MHRA 2012
- M3003: The Expression of Uncertainty and Confidence in Measurement: UKAS
- Retention and Storage of Pathological Records and Specimens - RCPATH, 2015, 5thEd
- Haemoglobinopathy Diagnosis 2nd Edition 2005: B.J Bain.
- Handbook of Transfusion Medicine: Fifth Edition 2014
- Dacie and Lewis Practical Haematology 12th Ed: S M Lewis, B J Bain, I Bates, M Laffan
- Blood Cells. A practical Guide 5th Edition 2015: B J Bain.
- NHSGGC: Confidentiality and Data Protection Policy
- NHSGGC: Confidentiality Policy
- NHS Code of Practice on Protecting Patient Confidentiality



Appendix 1

Changes from Previous Version

Update B12 reference ranges
Add telephone number of Transfusion practitioner

